

Presto™ Mini RNA Bacteria Kit

RBB004, RBBD004 (4 Preparation Sample Kit)

RBB050, RBBD050 (50 Preparation Kit)

RBB100, RBBD100 (100 Preparation Kit)

RBB300, RBBD300 (300 Preparation Kit)

Advantages

Sample: Gram (+) positive and Gram (-) negative bacterial cells

Yield: up to 60 µg of RNA (1 x 10⁹ *Escherichia coli*: 40-45 µg, 1 x 10⁹ *Bacillus subtilis*: 50-55 µg)

Convenient: includes Lysozyme and Bacteria Lysis Buffer

Format: certified DNase and RNase-free spin columns

Time: within 20 minutes

Elution Volume: 50-100 µl

Kit Storage: dry at room temperature (15-25°C), DNase I and Lysozyme are shipped at room temperature and should be stored at -20°C for extended periods

Table of Contents

Introduction.....	2
Quality Control.....	2
Kit Components.....	2
Safety Measures.....	3
Quick Protocol Diagram.....	3
Protocol Procedure.....	4
Troubleshooting.....	6
Test Data.....	6
Related Products.....	7

Introduction

The Presto™ Mini RNA Bacteria Kit was designed for total RNA purification from Gram (-) negative and Gram (+) positive bacteria. The provided Lysozyme and Bacteria Lysis Buffer will efficiently lyse bacterial cell walls consisting of the peptidoglycan layer. Detergents and chaotropic salt are used to further lyse cells and inactivate RNase while RNA is bound by the glass fiber matrix of the RNA spin column. Once any contaminants have been removed, using the Wash Buffer (containing ethanol), the purified total RNA is eluted by RNase-free Water and is ready for use in a variety of subsequent reactions.

Quality Control

The quality of the Presto™ Mini RNA Bacteria Kit is tested on a lot-to-lot basis by isolating RNA from *Escherichia coli* (1×10^9) culture (OD₆₀₀=1.3, 1 ml) harvested by centrifugation at 16,000 x g for 1 minute. 10 µl from a 50 µl eluate of purified RNA is analyzed by electrophoresis on a 0.8% agarose gel.

Kit Components

Component	RBB004 RBB004	RBB050 RBB050	RBB100 RBB100	RBB300 RBB300
Bacteria Lysis Buffer	1.5 ml	15 ml	30 ml	75 ml
Lysozyme ¹	8 mg	110 mg	250 mg	610 mg
RB Buffer	2 ml	30 ml	60 ml	130 ml
DNase I ² (2U/µl) (RBB004/050/100/300 Only)	20 µl	275 µl	550 µl	550 µl x 3
DNase I Reaction Buffer (RBB004/050/100/300 Only)	200 µl	2.5 ml	5 ml	15 ml
W1 Buffer	2 ml	30 ml	50 ml	130 ml
Wash Buffer ³ (Add Ethanol)	1.5 ml (6 ml)	25 ml (100 ml)	25 ml + 12.5 ml (100 ml) (50 ml)	50 ml x 2 (200 ml x 2)
RNase-free Water	1 ml	6 ml	15 ml	30 ml
RB Columns	4	50	100	300
2 ml Collection Tubes	8	100	200	600

^{1,2}Lysozyme and DNase I are shipped at room temperature and should be stored at -20°C for extended periods after receiving the kit.

³Add absolute ethanol (see the bottle label for volume) to Wash Buffer then mix by shaking for a few seconds. Check the box on the bottle. Be sure and close the bottle tightly after each use to avoid ethanol evaporation. The additional **Wash Buffer x 12.5 ml** is **only** included in **RBB100**.

Steps to prevent RNase contamination

1. During operation, always wear a lab coat, disposable gloves, protective goggles and (anti-fog) procedure mask.
2. Disposable plasticware and automatic pipettes should be sterile (RNase-free) and used only for RNA procedures.
3. Non-disposable glassware or plasticware should also be sterile (RNase-free).




During the procedure, always wear a lab coat, disposable gloves, and protective goggles.


Quick Protocol Diagram



Cell lysis of bacteria samples



RNA binding to membrane while contaminants remain suspended



Wash (removal of contaminants while RNA remains bound to membrane)



Elution of pure total RNA which is ready for subsequent reactions

Presto™ Mini RNA Bacteria Kit Protocol

Please read the entire instruction manual prior to starting the Protocol Procedure.

IMPORTANT BEFORE USE!

Add absolute ethanol (see the bottle label for volume) to Wash Buffer then mix by shaking for a few seconds. Check the box on the bottle. Be sure and close the bottle tightly after each use to avoid ethanol evaporation.

DNA Removal Options: For DNA-free RNA perform either option 1 (following RNA Binding) or option 2 (following RNA Elution).

Additional Requirements

absolute ethanol, ddH₂O (RNase-free and DNase-free) to prepare 70% ethanol, microcentrifuge tubes (RNase-free), pipette tips (RNase-free), β-mercaptoethanol, 15 ml centrifuge tube (RNase-free)

Bacteria Protocol Procedure

1. Sample Preparation

Transfer **bacterial cells (up to 1×10^9)** to a **1.5 ml microcentrifuge tube (RNase-free)**. Centrifuge for 1 minute at 14-16,000 x g then remove the supernatant completely. Transfer required volume of **Bacteria Lysis Buffer (200 μ l/sample)** to a **15 ml centrifuge tube (RNase-free)**. Add **Lysozyme (2 mg/200 μ l)** to **Bacteria Lysis Buffer (in the 15 ml centrifuge tube)** and vortex to completely dissolve the Lysozyme. Transfer **200 μ l of Bacteria Lysis Buffer (make sure Lysozyme was added)** to the sample in the 1.5 ml microcentrifuge tube then re-suspend the pellet by pipetting. Incubate at room temperature for 10 minutes. During incubation, invert the tube every 2-3 minutes.

2. Cell Lysis

Add **300 μ l of RB Buffer and 3 μ l β -mercaptoethanol (or 6 μ l of freshly prepared 2M Dithiothreitol in RNase Free Water)** and vortex. Incubate at room temperature for 5 minutes then centrifuge at 14-16,000 x g for 2 minutes. Transfer the supernatant to a new 1.5 ml microcentrifuge tube (RNase-free).

3. RNA Binding

Add **500 μ l of 70% ethanol to the lysate** and pipette immediately. Place a **RB Column** in a 2 ml Collection Tube. **Transfer 500 μ l of the mixture to the RB Column**. Centrifuge at 14-16,000 x g for 1 minute then discard the flow-through. Transfer the remaining mixture to the same **RB Column** and centrifuge at 14-16,000 x g for 1 minute. Discard the flow-through and place the RB Column in a new 2 ml Collection Tube.

Optional Step 1: In Column DNase I Digestion

The amount of DNA contamination is significantly reduced following In Column DNase I Digestion. However, traces of residual DNA may be detected in very sensitive applications. In this situation, please perform Optional Step 2: DNA Digestion In Solution instead to efficiently remove trace amounts of DNA. Standard DNase buffers are incompatible with In Column DNase I Digestion and may affect RNA integrity and reduce yield.

1. Add 400 μ l of Wash Buffer (make sure ethanol was added) to the RB Column then centrifuge at 14-16,000 x g for 30 seconds.
2. Discard the flow-through and place the RB Column back in the 2 ml Collection Tube.
3. Prepare DNase I solution in a 1.5 ml microcentrifuge tube (RNase-free) as follows:

DNase I	5 μ l (2 U/ μ l)
DNase I Reaction Buffer	45 μ l
Total Volume	50 μ l

4. Gently pipette DNase I solution to mix (DO NOT vortex) then add DNase I solution (50 μ l) into the CENTER of the RB column matrix.
5. Incubate the column for 15 minutes at room temperature (20-30°C) then proceed with Step 4 RNA Wash.

4. RNA Wash

Add **400 µl of W1 Buffer to the RB Column** then centrifuge at 14-16,000 x g for 1 minute. Discard the flow-through and place the **RB Column** back in the 2 ml Collection Tube. Add **600 µl of Wash Buffer (make sure ethanol was added)** into the **RB Column**. Centrifuge at 14-16,000 x g for 1 minute. Discard the flow-through then place the **RB Column** back in the 2 ml Collection Tube. Add **600 µl of Wash Buffer (make sure ethanol was added)** into the **RB Column**. Centrifuge at 14-16,000 x g for 1 minute. Discard the flow-through then place the **RB Column** back in the 2 ml Collection Tube. Centrifuge at 14-16,000 x g for 3 minutes to dry the column matrix.

5. RNA Elution

Place the **dried RB Column** in a clean 1.5 ml microcentrifuge tube (RNase-free). Add **50 µl of RNase-free Water** into the **CENTER** of the column matrix. Let stand for at least 3 minutes to ensure the **RNase-free Water** is absorbed by the matrix. Centrifuge at 14-16,000 x g for 1 minute to elute the purified RNA.

Optional Step 2: DNA Digestion In Solution

1. Prepare DNase I reaction in a 1.5 ml microcentrifuge tube (RNase-free) as follows:

RNA in RNase-free Water	1-40 µl
DNase I	0.5 µl/µg RNA
DNase I Reaction Buffer	5 µl
RNase-free Water	Add to final volume = 50 µl
Total Volume	50 µl

2. Gently pipette the DNase I reaction solution to mix (DO NOT vortex) then incubate the microcentrifuge tube at 37°C for 15-30 minutes.

3. Stop the reaction by adding 1 µl of 20 mM EGTA (pH=8.0) then incubate the microcentrifuge tube at 65°C for 10 minutes.

NOTE: DNase I Reaction Buffer may cause aberrant migration or smearing of RNA on gels. If analyzing RNA by gel electrophoresis, repurify the RNA sample by using the Geneaid™ RNA Cleanup Kit instead of stopping the reaction with EGTA.

Troubleshooting



Low Yield

Clogged Column.

Reduce the amount of starting material or separate it into multiple tubes. Centrifugation temperature must be between 20°C to 25°C. Bacteria cells were not completely homogenized. Make sure Lysozyme was added to Bacteria Lysis Buffer immediately prior to use.

Residual Ethanol Contamination.

Following the wash step, dry the RB Column with additional centrifugation at 14-16,000 x g for 5 minutes.

RNA Degradation.

The harvested sample should be stabilized immediately prior to use. Disposable plasticware and automatic pipettes should be sterile (RNase-free) and used only for RNA procedures. Non-disposable glassware or plasticware should also be sterile (RNase-free).

Presto™ Mini RNA Bacteria Kit Functional Test Data




Figure 1. RNA was extracted using the Presto™ Mini RNA Bacteria Kit. An *Escherichia coli* (1×10^9) culture (OD₆₀₀=1.3, 1 ml) was harvested by centrifugation at 16,000 x g for 1 minute. 10 µl from a 50 µl eluate of purified RNA was analyzed by electrophoresis on a 0.8% agarose gel.

Test	RNA Yield	260/280	260/230
1	41.56 µg	2.14	2.35
2	40.87 µg	2.15	2.32

Related RNA/DNA Extraction Products

RNA Purification		
Product	Package Size	Catalogue Number
Total RNA Mini Kit (Blood/Cultured Cell)	50/100/300 preps	RB050/100/300
Total RNA Mini Kit (Tissue)	50/100/300 preps	RT050/100/300
Total RNA Mini Kit (Plant)	50/100/300 preps	RP050/100/300
Presto™ Mini RNA Bacteria Kit	50/100/300 preps	RBB050/100/300
Presto™ Mini RNA Yeast Kit	50/100/300 preps	RY050/100/300
miRNA Isolation Kit	50/100 preps	RMI050/100
GENEzol™ Reagent	50/100/200 rxns	GZR050/100/200
GENEzol™ TriRNA Bacteria Kit	50/100 rxns	GZB050/100
GENEzol™ TriRNA Pure Kit	50/100/200 preps	GZX050/100/200
TriRNA Pure Kit	50/100/200 preps	TRP050/100/200
RNA Cleanup Kit	50/100 preps	PRO50/100
Virus DNA/RNA Purification		
Product	Package Size	Catalogue Number
Plant Virus RNA Kit	50/100 preps	PVR050/100
Viral Nucleic Acid Extraction Kit II	50/100/300 preps	VR050/100/300
Viral Nucleic Acid Extraction Kit III	50/100/300 preps	VI050/100/300
Genomic DNA Extraction		
Product	Package Size	Catalogue Number
Genomic DNA Mini Kit (Blood/Cultured Cell)	100/300 preps	GB100/300
Genomic DNA Maxi Kit (Blood/Cultured Cell)	10/25 preps	GDM010/25
Genomic DNA Mini Kit (Tissue)	50/100/300 preps	GT050/100/300
gSYNC™ DNA Extraction Kit	50/100/300 preps	GS050/100/300
Genomic DNA Mini Kit (Plant)	100 preps	GP100
Genomic DNA Maxi Kit (Plant)	10/25 preps	GPM10/25
GENEzol™ DNA Reagent Plant	100/200 rxns	GR100/200
Presto™ Mini gDNA Yeast Kit	100/300 preps	GBY100/300
Presto™ Mini gDNA Bacteria Kit	100/300 preps	GBB100/101/300/301
Genieus™ Micro DNA Extraction Kit	100/300 preps	GMB100/300
Presto™ Buccal Swab gDNA Extraction Kit	100/300 preps	GSK100/300
Presto™ 96 Well Blood Genomic DNA Extraction Kit	4/10 x 96 preps	96GBP04/10
Presto™ 96 Well Plant Genomic DNA Extraction Kit	4/10 x 96 preps	96GPP04/10
DNA RNA Purification		
Product	Package Size	Catalogue Number
Presto™ DNA/RNA Extraction Kit	50/100 preps	DR050/100
Presto™ DNA/RNA/Protein Extraction Kit	50/100 preps	DRP050/100

For additional product information please visit www.geneaid.com. Thank you!



Geneaid

