

1 **The Foster method: Rapid and non-invasive detection of clinically significant**  
2 **American Foulbrood disease levels using eDNA sampling and a dual-target**  
3 **qPCR assay, with its potential for other hive pathogens.**

4

5 John F. Mackay<sup>1\*</sup>, Rebecca E. Hewett<sup>1</sup>, Noa T. Smith<sup>1</sup> Tammy L. Waters<sup>1,3</sup>, John S.  
6 Scandrett<sup>2</sup>

7 <sup>1</sup> dnature diagnostics & research Ltd, 60 Carnarvon St., Gisborne, New Zealand

8 <sup>2</sup> Scandrett Rural Ltd, Invercargill, New Zealand

9 <sup>3</sup> current address: Environmental Science & Research, Porirua, New Zealand

10

11 \*Email: [john@dnature.co.nz](mailto:john@dnature.co.nz) Tel: +64 6 8633303

12

## 13 **Abstract**

14 Clinical signs of American Foulbrood can be difficult to diagnose and thus disease is  
15 missed and leads to further spread. Diagnosis is centred on the beekeeper's skill in  
16 recognising clinical symptoms – a highly subjective and time-consuming activity.  
17 Previous testing methods have relied on sampling that necessitates dismantling the  
18 hive and/or requires multiple visits to retrieve passive samples. The Foster method is  
19 a novel environmental DNA sampling method using entrance swabs together with a  
20 dual-target qPCR for AFB. The quantification data generated can be used to detect  
21 hives with clinically significant infections, even before visual symptoms are apparent.  
22 Such a method will be applicable to other bee pathogens and incursion pests.

23

24 **Keywords:** honey bee, American foulbrood, entrance, qPCR, diagnosis, swab,  
25 quantification, non-invasive.

## 26 **Importance**

27 Discovery of the disease American foulbrood typically means the destruction of the  
28 bees and hive by burning. This discovery is typically by visual examination of capped  
29 brood by the beekeeper - a subjective skill that means the disease is being missed  
30 or not recognised. It is a time-consuming and exacting process to inspect hives for  
31 AFB. Here we present a rapid sampling method that does not require opening/  
32 dismantling the hive, in conjunction with a dual target quantitative PCR assay for the  
33 bacteria responsible, *Paenibacillus larvae*. Using the resulting quantitative data,  
34 hives presenting clinical symptoms or likely to become clinical visually can be  
35 determined and the hives dealt with appropriately before further spread occurs.

36

## 37 **Introduction**

38 American Foulbrood (AFB) is one of the most destructive diseases in honeybees  
39 (*Apis mellifera*) and is caused by spores of the Gram positive bacteria *Paenibacillus*  
40 *larvae* infecting bees during their larval stage. These spores are extremely hardy;  
41 being resistant to heat, caustic, and other chemical treatments with the ability to  
42 remain infectious for over 30 years (Genersch, 2010). Only a few spores are  
43 required to cause an infection in a new larva and yet a single infected larvae can  
44 generate many millions of spores - leading to rapid spread and decline in a bee  
45 colony. Once the colony has weakened or died then it may be 'robbed' by bees from  
46 neighbouring hives where contaminated honey is brought back to a neighbouring  
47 hive, which leads to further infections. Infection is also spread unknowingly by the  
48 beekeeper (Fries and Camazine, 2001), through the introduction of hive frames  
49 carrying an undetected infection into a new colony of bees or the feeding of  
50 contaminated honey to colonies.

51 In New Zealand, hive numbers have more than doubled since 2013  
52 ([www.afb.org.nz](http://www.afb.org.nz)). Similarly the number of beekeepers has also increased – primarily  
53 due to hobby and conservation efforts (several hives per beekeeper) but also in the  
54 commercial sector (hundreds or thousands of hives per operation). Much of the  
55 growth in hive and commercial beekeeping has been due to the increased demand  
56 for high-value New Zealand manuka honey. The increase in hive numbers (and  
57 densities) combined with newer beekeepers, who are less likely to be skilled in  
58 identifying the clinical signs of AFB, has led to an increase in the percentage of hives  
59 reported by beekeepers as infected from 0.21% in 2015 to 0.32% in 2017  
60 ([www.afb.org.nz](http://www.afb.org.nz)). Since this time, the reported percentage has remained static,  
61 while hive numbers have continued to increase. ([www.afb.org.nz](http://www.afb.org.nz)).

62 The diagnosis of AFB is based primarily on visual symptoms – typically unusual,  
63 capped brood cells on a beehive frame that warrant investigation through a ‘roping’  
64 field test. In a hive with three to seven frames of brood or more, this means typically  
65 thousands of brood cells to examine visually per hive. Other diseases can affect  
66 brood cells meaning that examining suspicious cells for clinical AFB signs is  
67 laborious and prone to disease being missed. Clinical signs can vary based on the  
68 AFB genotype present, with ERIC II strains exhibiting reduced clinical symptoms  
69 (Genersch, 2010). In addition, hives must be dismantled, and the colony disrupted in  
70 order to inspect for AFB.

71 Confirmation for AFB in beehive materials has been traditionally performed by  
72 culture, using selective media, and induced germination of spores. However this is a  
73 variable process (Forsgren, 2008) and may take up to a week for results. PCR and  
74 qPCR (quantitative PCR) have been demonstrated as effective tools for the  
75 detection of AFB, with qPCR being the preferred diagnostic tool in recent years due  
76 its faster time to results and lower chance for amplicon contamination. However most  
77 applications have not used the quantitative data generated; rather using the tool for  
78 rapid confirmation of culture isolates (Kňazovická et al, 2011). The quantification  
79 aspect is important as low levels of AFB are often not clinically relevant (Bassi et al.,  
80 2018 , Pernal and Melathopoulos, 2006) and such hives may never develop clinical  
81 symptoms. Much of this low level may be due to the hygienic behaviour of bees to  
82 reduce overt infections (Spivak and Reuter, 2001).

83 The vast majority of molecular publications relating to AFB to date have relied on  
84 either conventional PCR or SYBR Green-based qPCR and the sole use of the 16S  
85 gene due to its high sensitivity as a multicopy target (Dobbelaere *et al.*, 2001; Piccini  
86 *et al.*, 2002; ). However this sensitivity can be at the expense of specificity, as

87 evidenced by the initial design (Martinez, 2010) and subsequent re-evaluation/re-  
88 design of qPCR primers to improve specificity (Rossi *et al.*, 2018). Sensitivity for  
89 bacterial targets can also be compromised by deletions of target sequences or  
90 sequence polymorphisms which affect primer or probe binding (Dahlberg *et al.*,  
91 2018; Johansen *et al.*, 2019; Xiu *et al.*, 2021) and many microbiological qPCR tests  
92 now utilise two independent targets to enhance both sensitivity and specificity, such  
93 as many of the commercial qPCR tests for chlamydia. Hydrolysis probe (“TaqMan”)  
94 qPCR assays permit quantification of each target by selected fluorescent  
95 wavelengths while also allowing the use of an internal control in an additional  
96 detection channel for confirmation of suitable quality DNA present in the reaction.  
97 These internal controls may improve the sensitivity of previous studies whereby the  
98 presence of inhibitors possibly prevented the amplification of AFB targets. (Forsgren  
99 and Laugen, 2013)

100 We sought to use a multiplex qPCR assay for AFB and evaluate a new and rapid  
101 environmental DNA (eDNA) sampling method for the detection of hives having - or at  
102 risk of developing - clinical AFB infections. Previously AFB has been detected in  
103 debris from a hive baseboard (Bassi *et al.*, 2018) but this study required installing  
104 collection sheets, necessitating dismantling the hive twice as well as repeat visits.  
105 The ability to detect relevant levels of AFB from a single sample taken from the  
106 entrance of a hive would permit far more rapid sampling of hives and allow infected  
107 hives to be dealt with more quickly (Lyll *et al.*, 2019). In this study, we evaluated  
108 and compared the AFB levels among bees, the entire baseboard, and the entrance  
109 region of the baseboard.

## 111 **Methods**

112 Quarantining hives in an apiary upon discovery of a clinical AFB hive is a common  
113 international practice, in order to minimise spread of the disease. From five  
114 quarantine apiaries around New Zealand (East Coast, Wellington, Wairarapa and  
115 South Canterbury regions) multiple hives were sampled two-monthly from late 2018  
116 to early 2020. In addition, samples from visually clinical hives and neighbouring hives  
117 were collected by local apiary inspectors for testing during this time. Samples  
118 included nurse bees, swabs from the hive entrance and swabs from over the entire  
119 baseboard. Subsequent samples comprised of comparisons between bees and  
120 solely the hive entrance. Bees were collected from the brood frames using 50 mL  
121 sterile containers and frozen to euthanise the bees. Sterile foam-tipped swabs  
122 (Puritan) were moistened in sterile water before swabbing either the whole  
123 baseboard or just through and across the hive entrance for three to five seconds.  
124 Instructions were provided to participating beekeepers to rotate the swabs during  
125 collection to ensure all swab surfaces could collect hive material. Swab heads were  
126 then snapped off into a 2 mL microcentrifuge tube, capped and sent to the laboratory  
127 with the bee samples, at ambient temperature by overnight courier. Once in the  
128 laboratory, bee samples were frozen at -20°C and swabs were stored at ambient  
129 temperature until DNA extraction was performed.

130 Genomic DNA extraction from bees and associated bacteria was performed from  
131 each sample containing 10 bees. DNA was extracted using the Bee Pathogen  
132 DNA/RNA Extraction Kit (dnature diagnostics & research Ltd, Gisborne, New  
133 Zealand) in conjunction with beadbeating. 10 bees were used for the practical  
134 purpose of ease of sample handling and throughput. In short, 3 mL DXL lysis buffer  
135 supplied in the kit was added to 10 bees in a 5 mL tube, containing a mixture of 0.5

136 mm (~2.4 g) and 2.3 mm zirconia silica (~1 g) beadbeating beads (BioSpec  
137 Products, Bartlesville, USA). The bees were homogenised in a Mini Beadbeater 16  
138 instrument (BioSpec Products, Bartlesville, USA) for 3 minutes. Tubes were then  
139 incubated at 65°C for 10 minutes with manual inversion every 3 minutes. 1 mL of  
140 the homogenate was transferred to a 2 mL microcentrifuge tube and centrifuged at  
141 15,000 x g for 5 minutes. 500 µL of supernatant was transferred to a new  
142 microcentrifuge tube containing 450 µL AD buffer and mixed well. The solution was  
143 then applied to a nucleic acid binding column from the kit (in two passes) and  
144 centrifuged through before successive washing steps and elution of nucleic acids  
145 from the column in 50 µL elution buffer.

146 A similar process was used for swabs: they were placed in a screwcap 2 mL tubes  
147 with 0.5mm zirconia silica beads (~0.8g) and 2.3 mm zirconia silica beads (0.3g).  
148 800 µL DXL buffer was added to each tube and the swab material homogenised in  
149 the Mini Beadbeater 16 for 3 minutes. The tubes were then incubated at 65°C with  
150 shaking for 10 minutes, before centrifuging and processing 500 µL of the resulting  
151 supernatant as for the bees. For pooled applications, swabs were added to 500 µL  
152 DXL buffer in a 2 mL microcentrifuge tube and vortexed for 10 seconds to dislodge  
153 cells and bacteria off the swab. 60 µL of each eluate (up to twelve swabs) was added  
154 to a 2 mL beadbeating tube as for single swabs and processed as for single swabs  
155 from the beadbeating step.

156 Assessment of DNA quality and presence of potential inhibitors was performed by  
157 measurement (spectroscopy) on a DS-11 FX (DeNovix, Wilmington, USA) and  
158 OD260/280 measurements recorded. Dilutions of extracted DNA were also tested by  
159 qPCR to measure the expected Cq increases with increasing dilution and thus  
160 monitor any qPCR inhibition.

161 The AFBduo qPCR test (dnature diagnostics & research Ltd, Gisborne, NZ) uses two  
162 targets – a single copy gene for quantification purposes (ftsZ, a prokaryotic  
163 homologue to tubulin) and the multicopy 16S gene. The ftsZ probe is detected in the  
164 FAM channel and the 16S target in the HEX channel. The test also includes a  
165 simultaneous internal control (ROX channel) for an animal mitochondrial marker, to  
166 confirm validity of results (in the cases where AFB was not detected) by way of  
167 successful internal control amplification.

168 Each AFBduo qPCR reaction was performed in a 10 µL volume and consisted of 5.5  
169 µL PCR grade water, 0.5 µL of the 20X Oligo Mix and 2 µL of the 5X Mastermix (all  
170 reagents supplied in the AFBduo real-time PCR kit, dnature). 2 µL genomic DNA,  
171 positive control DNA or sterile water (for no template controls) was added per  
172 reaction. The amplifications were performed on the Mic qPCR Cyclor (BioMolecular  
173 Systems, Australia) or the Eco qPCR instrument (Illumina, San Diego, USA) and the  
174 cycling conditions comprised: denaturation at 95°C for 2 minutes followed by 40  
175 cycles of 95°C for 5 seconds and 60°C for 15 seconds. Data was acquired on the  
176 Green (ftsZ), Yellow (16S) and Orange (Internal control) channels of the Mic cyclor  
177 or FAM, HEX and ROX channels on the Eco instrument. Cq's (Cycle quantities) were  
178 automatically generated by the Mic software, using dynamic analysis.

179 A synthetic DNA standard (gBlock, IDT, Singapore) was utilised that incorporated  
180 both amplicon sequences. The synthesised standard was resuspended in TE buffer,  
181 pH8, quantified on a DS-11 FX spectrophotometer (DeNovix, Wilmington, USA) and  
182 its copy number calculated using the equation:

$$183 \text{ Copies} = \frac{\text{ng} * \text{Avogadro's constant copies/mole}}{\text{length of standard} * 660 \text{ g/mole} * 1 \times 10^9 \text{ ng/g}}$$

184

185 Quantification standards were generated from  $10^6$  copies per  $\mu\text{L}$  down to 2 copies  
186 per  $\mu\text{L}$ . For all dilutions, the diluent was a pooled DNA sample extracted from AFB-  
187 free bees. Dilutions of extracted DNA from AFB-positive samples were diluted in  
188 AFB-free bee DNA. These dilutions were used to compare the sensitivity among  
189 existing World Organisation for Animal Health (OIE) conventional PCR protocols  
190 (Dobelaere, 1999; Govan, 2003) and the AFBduo qPCR assay. Conventional PCRs  
191 used the same 2  $\mu\text{L}$  each DNA dilution in a 20  $\mu\text{L}$  PCR using HOT FIREPol  
192 polymerase, ready to load (Solis BioDyne, Tartu, Estonia) and 0.3  $\mu\text{M}$  each primer.  
193 Following the recommended thermalcycling conditions (Dobelaere, 1999; Govan,  
194 2003), the conventional PCR reactions were analysed by electrophoresis on a 1%  
195 agarose gel.

196 The standards and their resulting Cq's were also used to convert the Cq's resulting  
197 from bee and swab samples to spore levels using the amplification efficiency of the  
198 single copy *ftsZ* gene, the y-intercept, and the formula:

199 
$$\text{Spores per swab} / 10 \text{ bees} = 10^{(\text{Cq}_{\text{ftsZ}} - 39) / -3.48} \times 25$$

200

## 201 **Results**

202 Testing of the AFBduo assay on synthetic DNA standards showed high analytical  
203 sensitivity, with both target reactions in the assay able to reliably detect down to 4  
204 copies per 10  $\mu$ L reaction. The reaction targets amplified with high efficiency in the  
205 multiplex (ftsZ = 94% and efficiency for 16S = 106% ) as well as excellent linearity  
206 over the 7 orders of magnitude ( $r = 0.998$ ). On testing AFB-positive bee, honey, and  
207 hive material, the 16S target typically amplified 3-5 cycles earlier (i.e. lower Cq's)  
208 than the ftsZ target as expected, due to the multiple copies of the 16S sequence per  
209 AFB bacterium compared to the single copy ftsZ gene.

210 The DNA extraction method from bees generated high purity DNA as shown by  
211 measurements on the microvolume spectrophotometer with OD260/280  
212 measurements of between 1.8 and 1.9. Extracted DNA samples serially diluted in  
213 elution buffer showed the expected increase in Cq's with decreasing concentration,  
214 indicating no inhibitors were present in the samples following DNA extraction. DNA  
215 yields from swabs were too low to be measured on the spectrophotometer, however  
216 internal control Cq's for serial dilutions as for the bees, indicated the DNA was of  
217 suitable quality.

218 Upon extracting DNA from AFB positive material, it was serially diluted it in  
219 honeybee DNA and these dilutions used to compare the AFBduo assays with the  
220 conventional PCR tests described in the OIE protocols (Dobbelare, 1999; Govan,  
221 1999). The AFBduo qPCR assay demonstrated two log greater sensitivity than the  
222 conventional PCR assays (data not shown).

223 In this study, we evaluated and compared the AFB levels among bees, the entire  
224 baseboard, and the entrance region of the baseboard. In the first cohort of

225 quarantine beehives from multiple regions in NZ (beehives from quarantine apiaries  
226 i.e. an apiary that had contained a hive exhibiting clinical signs), 45 hives were  
227 sampled up to 6 times during the 2018-2019 seasons. The samples taken were  
228 nurse bees, swabs from the entire baseboard and entrance swabs. Among these  
229 hives, 8 were found to have clinical signs of AFB ('roping' larval remains) during  
230 sampling while another 6 were presumed to have AFB due to their condition and the  
231 high prevalence in the apiary. Examples of the progression of AFB levels in three of  
232 the colonies is shown in Table 1, where the resulting Cq data has been converted to  
233 spores per 10 bee or swab sample processed.

234 In a second cohort of hives from one beekeeper, 23 hives were tested in a  
235 quarantine apiary using bee and hive entrance swab samples. Four of these hives  
236 showed clinical signs of AFB. Levels of AFB in nurse bee samples were compared  
237 to the levels indicated by the entrance swabs. Once again, most hives showed no or  
238 low levels of AFB, while those with clinical AFB demonstrated much higher spores  
239 levels of AFB in both the bee and entrance swab samples as shown in Table 2.

240 A third cohort of hives came from a beekeeper with a high incidence of AFB that was  
241 introduced through the purchase of contaminated equipment into his operation. Here  
242 we performed initial trials on pooling swab eluates into single extractions to reduce  
243 time and costs. Swab eluates were pooled into groups of eight and later twelve, the  
244 DNA extracted from the pooled sample and tested for the presence of AFB DNA.  
245 The remnant samples from pools that tested positive for both markers, were then  
246 tested individually as described. The results (Table 3) show that that even in the  
247 pools of twelve hive samples, single hives with higher levels of AFB could be clearly  
248 detected within a pool, prompting testing of those hives contributing to the positive  
249 pool to identify potentially clinically-affected hive(s).

## 250 Discussion

251 The AFBduo assay provided high confidence in the results due to the integrated  
252 internal control and detection of two markers for AFB. As expected, the 16S marker  
253 had approximately 10 to 20-fold higher analytical sensitivity due to the multi-copy  
254 nature of this marker. However the variability in Cq difference between the two  
255 markers among samples indicated the 16S copy number likely varied among AFB  
256 isolates and thus could not be relied on for quantification (Dahllof et al., 2000). This  
257 variability in copy number for diagnostic targets has been noted in other diagnostic  
258 qPCR assays for AFB (Rossi et al., 2018; Dainat et al., 2018). Therefore the use of  
259 the single copy *ftsZ* gene was used for quantification purposes using a standard  
260 curve prepared with a quantified synthetic template diluted in AFB-negative bee  
261 DNA.

262 Three sample types were compared in the first cohort – bees from the brood nest  
263 (nurse bees), swabs from the whole baseboard surface and swabs through the  
264 entrance of the hive only. The ability to sample the hive through the entrance  
265 removed the need to dismantle and disrupt the hive as with the first two sample  
266 types (bees and whole baseboard swabbing). The levels of both AFB and internal  
267 control DNA seen on the swabs taken from the whole baseboard or just the entrance  
268 area of the baseboard were similar (Table 4) possibly due to the maximum amount  
269 able to be collected on a given swab. This showed dismantling the hive to swab the  
270 whole baseboard was not required, in order to detect the higher levels of AFB in  
271 clinically-infected colonies. While the spore levels varied among clinical hives, they  
272 were still easily differentiated from the very low or undetectable spore levels in non-  
273 infected hives. While spore levels never increased during the monitoring time for  
274 many of the hives or became undetectable in some low spore cases, some of the

275 hives showed increasing levels at each testing point before clinical signs were  
276 observed in the respective hive and the hive destroyed (Table 1).

277 Spore levels were the highest in the bees from the brood nest area of the hive and  
278 while other studies have employed up to 100 bees per sample for culturing, the use  
279 of 10 bees permitted easier and more convenient handling for DNA extraction.

280 Despite the lower spore amounts on swabs, the levels from the entrance swabs were  
281 still high enough to be differentiated from levels seen in hives not exhibiting any  
282 signs of AFB (Table 2). While most hives without clinical signs of AFB had low or  
283 undetectable levels of DNA detected, one hive (Hive 11) showed levels similar to two  
284 other hives with clinical AFB signs. Indeed, at the next inspection by the beekeeper  
285 (approximately two weeks after sampling), hive 11 showed clinical signs of AFB and  
286 was destroyed thus demonstrating the ability of qPCR to find infected hives that may  
287 have pre-clinical infections or have symptoms missed by visual inspection.

288 However these results also demonstrated the importance of knowing the history of  
289 the hives sampled. In two cases (hives 11 and 12 in Table 2), the bees showed high  
290 levels of AFB spores while the hiveware showed low or undetectable levels.

291 Subsequent investigation showed that the hiveware was new (< 2 weeks old at time  
292 of sampling) for both these hives and that the colonies were newly-established from  
293 an unknowingly infected parent colony. Therefore, while the bees carried high levels  
294 of AFB there had not been enough time for spores to be deposited around the hive  
295 entrance surfaces. Further research will be required to assess this accumulation rate  
296 of spores on the hive.

297 Given the ability to detect clinically relevant levels of AFB through this simple hive  
298 entrance swabbing method, it is highly likely that the same DNA sample can be used

299 to test for other bee pathogens. *Nosema ceranae* has been detected through the use  
300 hive debris and qPCR (Copley *et al.*, 2012) and showed good correlation of levels  
301 between bees and debris. Since the development of this current work, a report has  
302 used hive debris collected over the period of a week to estimate loading of European  
303 foulbrood (a disease not present in New Zealand currently). As with the results  
304 described here, the bees offered a higher sensitivity of pathogen detection, yet the  
305 hive debris results still provided predictive information as compared to non-infected  
306 hives (Biová *et al.*, 2021). Another application could for biosecurity surveillance for  
307 unwanted hive pests such as tracheal mites (*Acarapis woodi*) or small hive beetle  
308 (*Aethina tumida*). The use of the sampling technique here combined with a qPCR  
309 assay (e.g. Li *et al.*, 2018; Ward *et al.*, 2007) could provide rapid detection of any  
310 incursion to countries such as New Zealand where small hive beetle is not found, by  
311 providing faster sampling of hives than dismantling hives to extract hive debris for  
312 testing (Ward *et al.*, 2007). AFB has also been detected from winter debris using  
313 conventional PCR and thus quantification was not possible (Ryba *et al.*, 2009). The  
314 quantification provides an important risk element of the hive - as to whether visual  
315 signs are likely present at the time of sampling or likely to demonstrate these clinical  
316 signs in the near future. In this work, levels of AFB spores of approximately 20,000  
317 spores/swab or greater, were associated with either clinical or pre-clinical levels of  
318 the disease.

319 All the methods published to date have required either dismantling the hive to  
320 sample from the baseboard or repeated visits to the hive to insert and remove  
321 sampling sheets. In contrast, the rapid sampling method described here that we  
322 have named the Foster method is a fast, one-time sampling and quantification

323 method that provides clinically significant information as to the AFB status of the

324 hive.

325

326

## 327 References

- 328 Bassi S, Carpana E, Bergomi P, Galletti G. 2018. Detection and quantification of  
329 *Paenibacillus* larvae spores in samples of bees, honey, and hive debris as a tool for  
330 American foulbrood risk assessment. *Bulletin of Insectology* 71 (2): 235-241.
- 331 Biová J, Charrière J-D, Dostálková S, Škrabišová M, Petřivalský M, Bzdil J, Danihlík,  
332 J. 2021. *Melissococcus plutonius* Can Be Effectively and Economically Detected  
333 Using Hive Debris and Conventional PCR. *Insects* 12, no. 2: 150
- 334 Copley T, Giovenazzo P, Jabaji S. 2012. Detection of *Nosema apis* and *N. ceranae*  
335 in honeybee bottom scraps and frass in naturally infected hives. *Apidologie* 43, 753–  
336 760.
- 337 Dahlberg J, Hadad R, Elfving K, Larsson I, Isaksson J, Magnuson A, Fredlund H,  
338 Unemo M, Herrmann B. 2018. Ten years transmission of the new variant of  
339 *Chlamydia trachomatis* in Sweden: Prevalence of infections and associated  
340 complications. *Sex. Transm. Infect.*, 94, 100–104.
- 341 Dahllöf I, Baillie H, Kjelleberg S. rpoB-based microbial community analysis avoids  
342 limitations inherent in 16S rRNA gene intraspecies heterogeneity. *Appl Environ*  
343 *Microbiol.* 2000 Aug;66(8):3376-80.
- 344 Dainat B, Grossar D, Ecoffey B, Haldemann C. 2018. Triplex real-time PCR method  
345 for the qualitative detection of European and American foulbrood in honeybee. *J.*  
346 *Microbio. Methods.* 146:61-63.
- 347 Dahllöf I, Baillie H, Kjelleberg S. rpoB-based microbial community analysis avoids  
348 limitations inherent in 16S rRNA gene intraspecies heterogeneity. *Appl Environ*  
349 *Microbiol.* 2000;66(8):3376-3380.
- 350 Dobbelaere W, de Graaf DC, Peeters JE, Jacobs FJ. 2001. Development of a fast  
351 and reliable diagnostic method for American foulbrood disease (*Paenibacillus* larvae  
352 subsp. larvae) using a 16S rRNA gene based PCR. *Apidologie* 32, 363–370.
- 353 Forsgren E, Stevanovic J, Fries I. 2008. Variability in germination and in temperature  
354 and storage resistance among *Paenibacillus* larvae genotypes, *Vet. Microbiol.* 129,  
355 342–349.

- 356 Forsgren E, Laugen A. 2014. Prognostic value of using bee and hive debris samples  
357 for the detection of American foulbrood disease in honey bee colonies. *Apidologie*  
358 45, 10–20.
- 359 Fries I, Carmazine S. 2001. Implications of horizontal and vertical pathogen  
360 transmission for honey bee epidemiology. *Apidologie* 32:199–214
- 361 Genersch E. 2010. American foulbrood in honeybees and its causative agent,  
362 *Paenibacillus larvae*. *J Invertebr Pathol* 103:S10–S19
- 363 Govan V, Allsopp M, Davidson S. 1999. A PCR detection method for rapid  
364 identification of *Paenibacillus larvae*. *Appl. Environ. Microbiol.*, 65, 2243–2245.
- 365 Johansen T, Kløvstad H, Rykkvin R, Herrfurth-Erichsen E, Sorthe J, Njølstad G.  
366 2019. The 'Finnish new variant of *Chlamydia trachomatis*' escaping detection in the  
367 Aptima Combo 2 assay is widespread across Norway, June to August 2019. *Euro*  
368 *Surveill* 24: 1900592.
- 369 Kňazovická V, Miluchová M, Gábor M, Kačániová M, Melich M, Kročko M,  
370 Trakovická A. 2011. Using Real-time PCR for Identification of *Paenibacillus larvae*.  
371 *Animal Science and Biotechnologies*, 44 (1).
- 372 Li D, Waite D, Fan Q, George S, Semeraro L, Blacket M. 2018. Molecular detection  
373 of small hive beetle *Aethina tumida* Murray (Coleoptera: Nitidulidae): DNA barcoding  
374 and development of a real-time PCR assay. *Sci Rep* 8, 9623.
- 375 Lyall D, Hansbro P, Horvat J, Stanwell P. 2020. Quantitative Nondestructive  
376 Assessment of *Paenibacillus larvae* in *Apis mellifera* Hives. In: Ahram T., Karwowski  
377 W., Pickl S., Tair R. (eds) *Human Systems Engineering and Design II. IHSED 2019.*  
378 *Advances in Intelligent Systems and Computing*, vol 1026. Springer, Cham.  
379 [https://doi.org/10.1007/978-3-030-27928-8\\_88](https://doi.org/10.1007/978-3-030-27928-8_88).
- 380 Martínez J, Simon V, Gonzalez B, Conget P. 2010. A real-time PCR-based strategy  
381 for the detection of *Paenibacillus larvae* vegetative cells and spores to improve the  
382 diagnosis and the screening of American foulbrood. *Lett. Appl. Microbiol.*, 50, 603–  
383 610.
- 384 Pernal S, Melathopoulos A. 2006. Monitoring for American Foulbrood spores from  
385 honey and bee samples in Canada. *Apiacta*, 41, 99–109.

- 386 Piccini C; D'Alessandro B; Antúnez K; Zunino P. 2002. Detection of *Paenibacillus*  
387 *larvae* subspecies larvae spores in naturally infected bee larvae and artificially  
388 contaminated honey by PCR. World J. Microbiol. Biotechnol., 18, 761–765.
- 389 Rossi F, Amadoro C, Ruberto A, Ricchiuti L. 2018. Evaluation of quantitative PCR  
390 (qPCR) *Paenibacillus* larvae targeted assays and definition of optimal conditions for  
391 its detection/quantification in honey and hive debris. Insects, 9, 165.
- 392 Spivak M, Reuter G. 2001. Resistance to American foulbrood disease by honey bee  
393 colonies *Apis mellifera* bred for hygienic behavior. Apidologie 32:555–565.
- 394 Ward L, Brown M, Neumann P, Wilkins S, Pettis J, Boonham N. 2007. A DNA  
395 method for screening hive debris for the presence of small hive beetle (*Aethina*  
396 *tumida*). Apidologie 38, 272–280.
- 397 Xiu L, Li Y, Zhang C, Li Y, Zeng Y, Wang F, Peng J. 2021. A molecular screening  
398 assay to identify *Chlamydia trachomatis* and distinguish new variants of *C.*  
399 *trachomatis* from wild-type. Microb Biotechnol. Mar;14(2):668-676.
- 400 Ryba S, Titera D, Haklova M, Stopka P. 2009. A PCR method of detecting American  
401 foulbrood (*Paenibacillus larvae*) in winter beehive wax debris. Vet. Microbiol., 139,  
402 193–196.
- 403

404 Hive FL5

Date	14 Sept 2018	4 Dec 2018	16 Jan 2019	Clinical status
Bees (spores/10 bees)	-	24,448	779,266	Clinical signs seen 16 Jan after sampling and hive destroyed
Entrance (spores/swab)	ND	10,870	678,363	

405

406 Hive STF4

Date	22 Sept 2018	Clinical status
Bees (spores/10 bees)	2,591,937	Clinical signs seen early October 2018 and hive destroyed
Entrance (spores/swab)	4,281,312	

407

408 Hive RL02

Date	31 Oct 2018	27 Dec 2018	11 Mar 2019	23 Aug 2019	Clinical status
Bees (spores/10 bees)	ND	ND	ND	ND	No AFB seen
Entrance (spores/swab)	ND	ND	ND	ND	

409

410 ND Not Detected

411 - Test not performed

412

413 **Table 1:** AFB spore levels seen in bee samples and entrance swabs from three  
 414 quarantine hives examples followed over time. Total number of hives followed over  
 415 time was thirty.

	Clinical signs of AFB?	Bees	Entrance Swab
		Spores/10 bees	Spores/swab
<b>MLHive 1</b>	<b>Yes</b>	<b>37,097,589</b>	<b>3,138,994</b>
<b>MLHive 2</b>	<b>Yes</b>	<b>514,061</b>	<b>24,008</b>
MLHive 3	No	ND	ND
MLHive 4	No	387	1,107
MLHive 5	No	539	377
MLHive 6	No	ND	ND
MLHive 7	No	80	ND
MLHive 8	No	210	ND
MLHive 9	No	205	ND
<b>MLHive 10</b>	<b>Yes</b>	<b>55,866,177</b>	<b>962,621</b>
MLHive 11	No**	9,518,751	ND*
<b>MLHive 12</b>	<b>Yes</b>	<b>12,561,091</b>	<b>890*</b>
MLHive 13	No	<10	<10
MLHive 14	No	ND	<10
MLHive 15	No	2049	1,318
MLHive 16	No	ND	861
MLHive 17	No	ND	326
MLHive 18	No	ND	1,092
MLHive 19	No	97	207
MLHive 20	No	ND	11,235
MLHive 21	No	ND	326
MLHive 22	No	ND	201
MLHive 23	No	943,739	2,365

416

417 **Table 2:** Testing an apiary comparing bee and entrance swab samples. The spore  
418 levels marked with \* indicate low level hive entrance spore on colonies recently  
419 placed into new hives (hives 11 and 12 were started from hive 10 splits). While at  
420 the time of sampling hive 11 showed no AFB symptoms (\*\*), it developed clinical

421 symptoms 2 weeks later and was destroyed.

422 ND Not detected

423

	Pooled result (12) ftsZ spore levels	Individual Swab Result ftsZ spore levels (per swab)	Clinical Signs of AFB (following testing)
Hive SB216	<b>15,630</b>	Not detected	No
Hive SB217		Not detected	No
Hive SB218		Not detected	No
Hive SB219		<b>165,103</b>	<b>Yes</b>
Hive SB220		Not detected	No
Hive SB221		Not detected	No
Hive SB222		Not detected	No
Hive SB250		Not detected	No
Hive SB251		Not detected	No
Hive SB252		Not detected	No
Hive SB253		Not detected	No
Hive SB254		Not detected	No
Hive SB255		<b>Not Detected</b>	-
Hive SB256	-		No
Hive SB257	-		No
Hive SB258	-		No
Hive SB259	-		No
Hive SB260	-		No
Hive SB261	-		No
Hive SB262	-		No
Hive SB263	-		No
Hive SB264	-		No
Hive SB265	-		No

424

425 **Table 3:** composite pooling of swabs to rapidly find clinically relevant levels in hives.

426 In this instance the first pool showed raised levels and the remnant swab eluates

427 were tested individually, whereupon elevated levels were found in hive SB219.

428 Subsequent hive inspections showed this hive to have clinical symptoms and it was

429 destroyed.

430 – test not performed

431

Hive	Location	Estimated Spore level (ftsZ target)	Hive status
ML Shearer	Baseboard	52 million	Clinical signs of AFB
	Entrance	25 million	
ML Ellis	Baseboard	3.6 million	Clinical signs of AFB
	Entrance	3.4 million	
FL 10	Baseboard	61	No AFB seen
	Entrance	335	

432

433 **Table 4:** Comparison of baseboard and entrance swabs showing the differentiation

434 of spore levels among two clinical hives and a hive without clinical symptoms.

435

436 **Conflict of interest statement**

437 JFM, REH and NTS are employees of dnature diagnostics & research Ltd., that  
438 developed the commercial AFBduo qPCR kit as well as the commercial bee DNA  
439 extraction kit method used in this study.

440

441 **Acknowledgements and memoriam**

442 This work was partly funded by Eastland Community Trust (now Trust Tairawhiti)  
443 (JFM, REH., NTS.,TLW) and also the Sustainable Food and Fibre Futures from  
444 Ministry of Primary Industries, Alpine Honey and NZ Honey Trust (JSS). We thank all  
445 the beekeepers who provided samples. We also thank Richard Hall (Ministry for  
446 Primary Industries) for early discussions and also Phil Lester (Wellington University)  
447 and Bart Janssen (Plant and Food Research) for comments on the manuscript. The  
448 method is named for NZ beekeeper Barry Foster - a long-time AFB educator,  
449 inspector, and former NZ AFB management agency board member. This paper is  
450 dedicated to the memories of Russell Poulter, a long-standing collaborator who  
451 passed away suddenly in January 2021 and beekeeper/AFB inspector John  
452 Maynard who passed away suddenly in February 2021.